

REFERRAL FORM

Name of Patient		
Appointment Date	Time	Diag. Code (ICD 9)
Referring Physician Signature		
Signs, Symptoms & Clinical Information		
Female Patient LMP		
<input type="radio"/> Routine Report Delivery <input type="radio"/> Fax Preliminary Report if Abnormal. Fax# _____		

Radiographic Exam

- Chest 1 View
- Chest PA & Lateral
- Ribs
 - Uni Bilateral Uni Ribs w/PA Chest
- Abdomen
 - One View Flat & Upright Acute Abdomen Series
- Skull AP & Lateral 4 Views
- Orbits
- Cervical Spine Complete w/Flex & Ext.
- Thoracic Spine AP & Lateral Complete
- Lumbar Spine AP & Lateral Complete
- Scoliosis AP Only Lat Only
- 77080 Bone Density
- Joints & Extremities (Designate) _____
- Other _____

Ultrasound Exam

- Complete Abdomen – Liver, GB, Pancreas, Kidneys
- Pelvis – Uterus, Ovaries (to include transabdominal and transvaginal)
- Kidney – Bladder
- Aorta
- Scrotum
- Thyroid
- Breast
- Extremity (popliteal cyst)
- Obstetrical Complete
- Obstetrical Limited
- Carotid Doppler
- Extremity Venous Doppler (DVT) Designate _____
- TIPS Evaluation
- Renal Transplants
- Groin r/o pseudo aneurysm
- Other _____

Mammography

- 77055 - Unilateral Mammogram (1 side)
- 77056 - Diagnostic Mammogram – To include ultrasound if indicated
- 77057 - Routine Mammogram – To include additional imaging and ultrasound if indicated
- 19290 - Needle Placement
- 77032 - Needle Loc. – Mammography
- 76098 – Breast Specimen
- 77032 – Needle Loc. Limited with Needle

PREPARATIONS

The following are routine adult preparations. If you feel that they are not indicated for your patient, please call for alternative instructions.

Clear Liquids include: Tea, coffee (no cream), soft drinks, boullion, jello (no fruit), apple juice, cranberry juice, ice popsicles.

Ultrasound:

Abdominal, Aorta, Biopsy and Tips Evaluation - Nothing to eat or drink after midnight.

Pelvic – Empty bladder at _____ (1-1/2 hour before your exam, start drinking 1 1/2 quarts of liquid. If you weigh over 150 pounds, drink 2 quarts.)

Do not empty bladder again until your exam is completed. Most pelvic and early obstetrical studies include an endovaginal ultrasound.

OB-Ultrasound – Follow pelvic prep for up to 20 weeks, after 20 weeks gestation, drink 1 quart only

Renal Ultrasound – Empty bladder 1 hour prior to exam. Drink 1 quart of noncarbonated liquid. Do not empty bladder again.

Renal Transplants – Same as Renal Ultrasound

PLEASE BRING THIS FORM WITH YOU FOR YOUR APPOINTMENT